



PubMed

National Library of Medicine
NLM

Entrez PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

Books

Search **PubMed**

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Nat Med. 2002 Feb;8(2):157-65.

Comment in:

• Nat Med. 2002 Feb;8(2):117-8.

**nature
medicine**

CCL27-CCR10 interactions regulate T cell-mediated skin inflammation.

Homey B, Alenius H, Muller A, Soto H, Bowman EP, Yuan W, McEvoy L, Lauerman AI, Assmann T, Bunemann E, Lehto M, Wolff H, Yen D, Marxhausen H, To W, Sedgwick J, Ruzicka T, Lehmann P, Zlotnik A.

DNAX Research Institute, Palo Alto, California, USA. berhard.homey@uni-duesseldorf.de

The skin-associated chemokine CCL27 (also called CTACK, ALP and ESKine) and its receptor CCR10 (GPR-2) mediate chemotactic responses of skin-homing T cells in vitro. Here we report that most skin-infiltrating lymphocytes in patients suffering from psoriasis, atopic or allergic-contact dermatitis express CCR10. Epidermal basal keratinocytes produced CCL27 protein that bound to extracellular matrix, mediated adhesion and was displayed on the surface of dermal endothelial cells. Tumor necrosis factor-alpha and interleukin-1beta induced CCL27 production whereas the glucocorticosteroid clobetasol propionate suppressed it. Circulating skin-homing CLA+ T cells, dermal microvascular endothelial cells and fibroblasts expressed CCR10 on their cell surface. In vivo, intracutaneous CCL27 injection attracted lymphocytes and, conversely, neutralization of CCL27-CCR10 interactions impaired lymphocyte recruitment to the skin leading to the suppression of allergen-induced skin inflammation. Together, these findings indicate that CCL27-CCR10 interactions have a pivotal role in T cell-mediated skin inflammation.

PMID: 11821900 [PubMed - indexed for MEDLINE]

Display **Abstract** Show: **20** **Sort** **Send to** **Text**

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services
Freedom of Information Act | Disclaimer





PubMed

National Library of Medicine
NLM

Entrez

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

Books

Search

PubMed

for

Limits

Preview/Index

History

Clipboard

Details

Go

Clear

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MESH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: J Dent Res. 2003 Aug;82(8):621-6.

Full text article at
jdr.sagepub.com

Differential injury responses in oral mucosal and cutaneous wounds.

Szpaderska AM, Zuckerman JD, DiPietro LA.

Burn and Shock Trauma Institute, Department of Surgery, Loyola University Medical Center, 2160 S. First Ave., Maywood, IL 60153, USA.

Oral mucosa heals faster than does skin, yet few studies have compared the repair at oral mucosal and cutaneous sites. To determine whether the privileged healing of oral injuries involves a differential inflammatory phase, we compared the inflammatory cell infiltrate and cytokine production in wounds of equivalent size in oral mucosa and skin. Significantly lower levels of macrophage, neutrophil, and T-cell infiltration were observed in oral vs. dermal wounds. RT-PCR analysis of inflammatory cytokine production demonstrated that oral wounds contained significantly less IL-6 and KC than did skin wounds. Similarly, the level of the pro-fibrotic cytokine TGF- β 1 was lower in mucosal than in skin wounds. No significant differences between skin and mucosal wounds were observed for the expression of the anti-inflammatory cytokine IL-10 and the TGF- β 1 modulators, fibromodulin and LTBP-1. These findings demonstrate that diminished inflammation is a key feature of the privileged repair of oral mucosa.

PMID: 1285847 [PubMed - indexed for MEDLINE]

Related Articles, Links

Display

Abstract

Show:

20

▼

Sort

▼

Send to

▼

Text

▼

▼

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer



Nov 18 2003 07:03:32

CCR10 Expression by Malignant Melanoma Cells: Implications for Tumor Growth and Metastasis

A. Müller¹, S. N. Wagner², E. P. Bowmann³, T. Ruzicka¹, A. Zlotnik⁴, and B. Homey¹.

¹Departments of Dermatology and Radiation Oncology, Heinrich-Heine University, Moorenstrasse 5, D-40225 Düsseldorf, Germany; ²Department of Dermatology, University of Vienna, Austria; ³DNAX Research Inc., Palo Alto, CA, USA; ⁴Eos Biotechnology, South San Francisco, CA, USA.

Malignant Melanoma shows aggressive primary tumor growth and metastasizes into local draining lymph nodes, lung, liver, brain and bone marrow. In contrast to other malignancies, malignant melanoma also shows a high frequency of skin metastases. Recently, we showed that chemokine receptors such as CXCR4 are involved in the development of organ-specific metastasis to lymph nodes and lung. Furthermore, we identified the novel skin-associated chemokine CCL27(CTACK, ALP, ESkin) and its receptor CCR10(GPR-2) which specifically mediate the recruitment of lymphocytes into the skin. In the present study, we demonstrate that malignant melanoma cell lines as well as primary malignant melanoma tumors express high levels of CCR10 mRNA. Notably, skin metastases of malignant melanoma show significantly increased CCR10 mRNA expression compared to primary tumors suggesting the selection for CCR10-high expressing malignant clones. Immunohistochemical analyses confirmed CCR10 expression by MelanA-positive melanoma cells. *In vitro*, CCR10 signaling mediated migration and invasion of malignant melanoma cells and significantly induced their proliferation. *In vivo*, neutralization of mCCL27 resulted in delayed primary tumor growth of human melanoma cells in a SCID mouse model. Taken together, our findings suggest that CCR10 expression by malignant melanoma cells plays a role in primary tumor growth and the development of skin metastases.



Meeting: Investigative Dermatology 2003 Meeting
Fontainebleau Hilton

Miami Beach, FL

April 30 - May 4, 2003